PRELIMINARY REPORT:

STATUS OF U.S. LAND REMOTE SENSING ARCHIVE

September 1989

Howard W. Warriner Landsat Archive Manager NOAA Landsat Operation Branch

TABLE OF CONTENTS

Introduction	1
Discussion	1
Procedure	2
Plots of Africa	3
Plots of South America	17

STATUS OF THE U.S. LAND REMOTE SENSING ARCHIVE, September 1989

INTRODUCTION

In 1989, an investigation was undertaken to determine and develop means that would display Landsat data listed in the National Satellite Land Remote Sensing Data Archive.

DISCUSSION

The outcome of the effort is a series of plots. As this is written, plots for the continents of Africa and South America for U.S. acquired Thematic Mapper data have been completed for 1983 through 1988. The 1989 data plot is, of course, incomplete at this time.

For each continent, there is a series of plots for all accession data by year and a summary for all years. In addition there is a series of plots depicting accessions with 30% or less cloud cover for all years and a summary of 30% or less cloud cover accessions for all years. Caution should be exercised in dealing with the cloud cover data plots. Generally the cloud cover value is determined by the Thematic Mapper Image Processing System (TIPS) machine. Historically, the TIPS cloud cover call was correct about 50% or less of the time.

Where large holidays were noted, such as the eastern coast of Brazil, EOSAT was informed, such that current data acquisitions could be obtained.

SUMMARY

Much more remains to be done. For example:

- a) The continents of North America, Antarctic, India, Europe, and Asia remain. I feel strongly that review of Antarctic data is required immediately for both MSS and TM ascending (day) and descending (night) data. I also believe that an ascending plot is required for the world.
- b) At this point, I believe that it would be instructive to modify the software routines to offer an option to incorporate a number in each frame which would depict the number of scenes available with the selection criteria used.
 - c) Other modifications such as; using color to depict satellite, season or other variables should be considered and incorporated.
 - d) Plots of international data; i.e., those accessions held by Landsat Ground Stations participating in the International Database at EDC could be accomplished.

PROCEDURE


To display the data available in the Archive, a series of procedures and some software programs were written. Briefly, the steps to accomplish these plots are as follows:

- a) Select area/sensor of interest and create the INORAC Inquiry (DI029) output tape.
- b) On a separate system, ingest the DI029 tape, reducing the 293 byte records to 128 bytes.
- c) Copy the smaller record to 5 1/4 or 3 1/2 inch IBM PC compatible floppies.
- d) Ingest the floppies with an IBM PC compatible machine (8086 through 80326).
- e) Apply a BASIC program to edit the 128 byte records, to select year, area, cloud cover, quality, etc. (DI29EDIT.BAS will do this.)
- f) Apply a BASIC program to compute the corner points for plotting with CAM. (MASTER.BAS will do this.)


g) Transfer the prepared CAM files to a machine licensed for CAM and execute.

This procedure is flexible enough to accommodate MSS, TM, and RBV data (no RBV testing was accomplished) along with ascending (night time) MSS and TM data for the U.S. and foreign holdings. However, time did not permit the testing of ascending data to determine if there are problems in plotting ascending data crossing zero degrees latitude and zero degrees longitude.


AFRICA - ALL TM DATA FOR 1984




AFRICA - ALL TM DATA FOR 1985




AFRICA - ALL TM DATA FOR 1986




AFRICA - ALL TM DATA FOR 1987




AFRICA - ALL TM DATA FOR 1988




AFRICA - ALL TM DATA FOR 1989




AFRICA - ALL TM DATA 1984 TO JUNE 1989




AFRICA - ALL TM DATA WITH 30% OR LESS CLOUD COVER FOR 1984




AFRICA - ALL TM DATA WITH 30% OR LESS CLOUD COVER FOR 1985




AFRICA - ALL TM DATA WITH 30% OR LESS CLOUD COVER FOR 1986




AFRICA - ALL TM DATA WITH 30% OR LESS CLOUD COVER FOR 1987




AFRICA - ALL TM DATA WITH 30% OR LESS CLOUD COVER FOR 1988




AFRICA - ALL TM DATA WITH 30% OR LESS CLOUD COVER TO JUNE 1989




AFRICA - ALL TM DATA WITH 30% OR LESS CLOUD COVER 1984 TO JUNE 1989







LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION MAP SCALE 1:45,000,000




LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION
MAP SCALE 1:45,000.000




LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION
MAP SCALE 1:45,000,000



LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION
MAP SCALE 1:45.000.000




LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION MAP SCALE 1:45,000,000




LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION MAP SCALE 1:45,000.000

SOUTH AMERICA TM DATA TO MARCH 1989




LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION MAP SCALE 1:45.000.000


SOUTH AMERICA TM DATA 1983 TO MARCH 1989




LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION
MAP SCALE 1:45,000,000




LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION
MAP SCALE 1:45,000,000




LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION
MAP SCALE 1:45,000,000




LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION
MAP SCALE 1:45.000.000



LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION
MAP SCALE 1:45.000.000




LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION MAP SCALE 1:45,000,000




LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION
MAP SCALE 1:45,000,000

SOUTH AMERICA TM DATA WITH 30 PERCENT OR LESS CLOUD COVER JANUARY TO MARCH 1989



LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION
MAP SCALE 1:45.000.000

SOUTH AMERICA TM DATA WITH 30 PERCENT OR LESS CLOUD COVER 1983 TO MARCH 1989



LAMBERT AZIMUTHAL EQUAL-AREA PROJECTION
MAP SCALE 1:45,000,000